



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2013

Intra-specific variation in social organisation by genetic variation, developmental plasticity, social flexibility or entirely extrinsic factors

Schradin, C

Abstract: Previously, it was widely believed that each species has a specific social organization, but we know now that many species show intraspecific variation in their social organization. Four different processes can lead to intraspecific variation in social organization: (i) genetic variation between individuals owing to local adaptation (between populations) or evolutionarily stable strategies within populations; (ii) developmental plasticity evolved in long-term (more than one generation) unpredictable and short-term (one generation) predictable environments, which is mediated by organizational physiological effects during early ontogeny; (iii) social flexibility evolved in highly unpredictable environments, which is mediated by activational physiological effects in adults; (iv) entirely extrinsic factors such as the death of a dominant breeder. Variation in social behaviour occurs between individuals in the case of genetic variation and developmental plasticity, but within individuals in the case of social flexibility. It is important to study intraspecific variation in social organization to understand the social systems of species because it reveals the mechanisms by which species can adapt to changing environments, offers a useful tool to study the ultimate and proximate causes of sociality, and is an interesting phenomenon by itself that needs scientific explanation.

DOI: <https://doi.org/10.1098/rstb.2012.0346>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-89698>

Journal Article

Accepted Version

Originally published at:

Schradin, C (2013). Intra-specific variation in social organisation by genetic variation, developmental plasticity, social flexibility or entirely extrinsic factors. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 368(1618):20120346.

DOI: <https://doi.org/10.1098/rstb.2012.0346>

Intra-specific variation in social organisation by genetic variation, developmental plasticity, social flexibility or entirely extrinsic factors

Schradin, C. 2013. Intra-specific variation in social organisation: genetic variation, developmental plasticity, social flexibility or purely extrinsic factors. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 1471-.

Carsten Schradin^{a,b,c,d}

^aUniversité de Strasbourg, IPHC-DEPE, 23 rue Becquerel 67087 Strasbourg, France

^bCNRS, UMR7178, 67087 Strasbourg, France

^cSchool of Animal, Plant and Environmental Sciences, University of the Witwatersrand, Johannesburg, South Africa

^dInstitute of Evolutionary Biology and Environmental Studies, University of Zurich, Switzerland.

carsten.schradin@iphc.cnrs.fr

Previously, it was widely believed that each species has a specific social organisation, but we know now that many species show intra-specific variation in their social organisation. Four different processes can lead to intra-specific variation in social organisation: 1. Genetic variation between individuals due to local adaptation (between populations) or evolutionary stable strategies within populations. 2. Developmental plasticity evolved in long-term (more than one generation) unpredictable and short-term (one generation) predictable environments and is mediated by organisational physiological effects during early ontogeny. 3. Social flexibility evolved in highly unpredictable environments and is mediated by activational physiological effects in adults. 4. Entirely extrinsic factors such as the death of a dominant breeder. Variation in social behaviour occurs between individuals in the case of genetic

variation and developmental plasticity, but within individuals in the case of social flexibility. Intra-specific variation in social organisation is important to study to understand the social systems of species because it reveals the mechanisms by which species can adapt to changing environments, it offers a useful tool to study the ultimate and proximate causes of sociality, and it is an interesting phenomenon by itself that needs scientific explanation.

Key words: social system; alternative reproductive tactics; population; frequency dependent selection; evolutionary stable strategy; striped mouse

1. INTRA-SPECIFIC VARIATION IN SOCIAL ORGANISATION

Understanding variation in social systems between species has long been one of the main aims in evolutionary biology (1-5). The social system of a species consists of the social organisation (the composition of groups), the social structure (describing who interacts with whom) and the mating system (6). Formerly, it was widely assumed that one species had one fixed form of social organisation. Divergence of single individuals from the species-specific pattern of social behaviour was often treated as noise in the data set. However, in the 1980s it was realized that variation in social systems occurs in many species, either between or within populations (7, 8).

Presently, the phylogenetic approach to understanding the evolution of different social systems is flourishing (9-11), but it requires each species to be categorised correctly with regard to its social organisation. However, apart from the standard categories (e.g. solitary, pair-living, multi male multi female species), there is no category for socially variable species. But to understand variation in social systems between species, it is important to understand the variation within species (7, 12). Here I focus on the social organisation, i.e. the composition of groups, because this is the parameter most easily measured in field studies, and because it influences the mating system, the social structure and thus the social system (6, 13).

Intra-specific variation in social organisation offers a unique opportunity to study the ultimate and proximate causes of social traits without any confounding phylogenetic effects (14, 15). Understanding the different mechanisms that underlie intra-specific variation in social organisation will also help us to understand how and whether species can adapt to changing environments, which would also inform conservation efforts. Finally, intra-specific variation in social organisation is itself an interesting phenomenon that needs scientific explanation both from the ultimate and the proximate perspective (7, 15).

While the occurrence of intra-specific variation in social organisation has received significant attention and acknowledgement recently, the focus has been on the associated ecological factors (13, 16, 17). Additionally, we need to understand the different proximate and evolutionary mechanisms that arise from these ecological factors. While many examples of population differences in social organisation exist (7), the underlying processes are typically not known. The aims of this review are to present an overview of the four processes that bring about intra-specific variation in social organisation (Fig. 1), and to critically evaluate the ecological (extrinsic) and physiological (intrinsic) factors that bring about this variation.

2. GENETIC VARIATION

Individuals of a species can differ genetically from each other in a way that influences their social behaviour (18, 19), such as their ability to form pair-bonds and social bonds, and their propensity to show parental care (20, 21). These individual differences in genotype influencing social behaviour could contribute to intra-specific variation in social organisation, both between and within populations.

If the environments of two populations of the same species differ in important aspects such as climate, availability of resources, pathogens, or the occurrence of other competing species, then natural and sexual selection can differ between populations, leading to local adaptations. For example, if the carrying capacity and thus population density differs between

populations due to environmental factors, different social tactics might evolve in the two populations. Another example would be the benefits of maternal / parental and bi-parental care (22), which can vary with changes in food availability, availability of mating partners and ambient temperature (23). Local adaptation might then result in genetic differences between populations regulating the expression of social behaviours, which can cause the social organisation of the two populations to differ. For example in prairie voles (*Microtus ochrogaster*) most individuals in some populations are pair-living, whereas the majority of individuals in other populations are solitary. Those two model types of social organisation might represent genetic adaptations to different environments (24) (but see also (25)).

Genetic variation could also contribute to variation in social organisation within populations. If the variation in social organisation occurs between generations, this could be due to frequency-dependent or fluctuating selection, such that different genotypes prevail at different times within the population. If variation in social organisation occurs at the same time, this might be due evolutionarily stable strategies (ESS) that are genetically determined (26). In this case, two or more genetically determined strategies exist within the population that have the same fitness and a higher fitness than any alternative strategy (1). Such ESS have been documented in the case of alternative reproductive tactics (ARTs). In side-blotched lizards (*Uta stansburiana*) three male tactics exist that are genetically different and maintained via frequency-dependent selection (27). Genetically determined ARTs that could represent ESS occur in several species of fish (28), lizards (27), in ruffs (*Philomachus pugnax*; (29), and in the isopod *Paracerceis sculpta* (30). In fire ants (*Solenopsis invicta*) the number of breeding queens per colony is determined by the single gene Gp-9, with one queen (monogynous social organisation) when only the B-allele is present, but multiple queens (polygynous social organisation) when the b-allele is present as well (31).

The existence of male ARTs alone might not change the social organisation unless females also show ARTs (15), but only in a few species do both sexes show ARTs. The extent

to which genetic differences can explain intra-specific variation in social organisation is not well understood, maybe because most studies focussed on male ARTs within one population, which is easier than making comparisons between populations. Still, genetic differences between populations due to local adaptation influencing social organisation might be common, and this could initiate ecological speciation (32).

3. PHENOTYPIC PLASTICITY

Phenotypic plasticity is the capacity of a specific genotype to produce different phenotypes (in behaviour, physiology or morphology) in response to different environmental conditions, including the social and pre-natal environment (33). Phenotypic plasticity evolves when (i) the environment of a population varies between or within generations, (ii) individuals can use reliable environmental cues indicating (iii) which phenotype has the highest fitness, meaning that (iv) different phenotypes have different fitness in different environments and (v) no phenotype exists that has the highest fitness in all environments (summarised by (34)).

Phenotypic plasticity can occur early in development and lead to permanent changes of the phenotype, or it can be reversible, which often occurs in adults. The first phenomenon has been called developmental plasticity, the second one phenotypic flexibility (35). Different physiological mechanisms exist for these two kinds of plasticity, which also suggest that different evolutionary forces are at work.

(a) Developmental plasticity

In the case of developmental plasticity, variation is due to environmental variation activating alternative developmental pathways of one genotype (35). Developmental plasticity is non-reversible, making it empirically difficult to differentiate it from genetic variation between individuals. This can be best achieved by experimental studies in captivity testing for an influence of the environment on the phenotype. If environmental manipulation has no influence

on the ratio of alternative phenotypes produced, genetic variation is likely to be the cause (36). If the environment significantly influences the resulting phenotype, developmental plasticity is likely to be at play (37). If in common garden experiments the observed differences in social organisation between populations disappear in the common garden, then developmental plasticity would be the explanation for the differences observed in nature.

Developmental plasticity is a response to the early environment within which an individual develops and in which it later grows up. For this, some cues of the current environment must reliably predict the future environment in which the same individual will reproduce, and this predictability must have occurred in the past when developmental plasticity evolved. As such, developmental plasticity enables the development of alternative adaptive phenotypes that have a higher fitness on average than any fixed phenotype. Typically, the environmental influence is during early development (pre- and / or postnatal), later affecting the phenotype of adult breeding individuals. This influence can occur before birth, for example due to the position in utero between male or between female siblings in rodents, and the resulting organisational effects due to the steroid hormones originating from the siblings (38). In rodents, maternal effects significantly influence stress response and social behaviour (39), which is an adaptive response to specific environments (40, 41). Developmental plasticity can also occur via environmentally induced changes in DNA methylation and as such represent epigenetic effects, causing phenotypic diversity (42).

If the environment of a population changes consistently, the development of the nervous system, and thus ultimately the social behaviour of this generation could significantly differ from that of other generations. Similarly, the environments of two different populations of the same species can differ significantly, inducing different developmental pathways, leading to intra-specific variation in social organisation. However, I am not aware of any empirical studies demonstrating the importance of developmental plasticity for intra-specific variation in

social organisation, which could well be due to researchers not addressing the necessary research questions.

(b) Social flexibility

Social flexibility describes the phenomenon that the social organisation of a species or population can change as a function of individuals reversibly changing their social tactics in response to short-term changes of the environment. Thus, social flexibility focusses on changes in the population that are a function of individuals of both sexes reversibly changing their reproductive and social tactics (15)). Individuals modify their interactions with other individuals (social structure), with whom they mate (mating system), and consequently the composition of groups (social organisation) and thus the entire social system of the population. In the case of social flexibility, both males and females must have alternative reproductive tactics based on a single strategy (all individuals have the same decision rules) (43). Ecological constraints are one of the most important factors for the evolution of social flexibility (15, 17). These constraints can vary between years, often due to changes in population density, such that in some years philopatry and the establishment of large groups are favoured, in other years dispersal and solitary breeding (5, 15, 17). Within a population, often two or more forms of social organisation might exist, e.g. solitary and group-living individuals, or monogamous and polygynous groups.

One problem with the term social flexibility is confusion arising from the fact that most behaviours are flexible (reversible). Many researchers in animal behaviour use the term “flexible” to emphasize this characteristic feature of social behaviour. As such, “flexible” is a vague term without scale (which behaviour is flexible, which is not?) and neither a scientific phenomenon nor a theoretical concept. In contrast, the value of the term “social flexibility” is that it provides a conceptual framework that enables us to study ultimate and proximate factors

of individual flexibility in social behaviour that leads to intra-specific variation in social organisation (15).

Interestingly, species that are flexible in their social behaviour might not show social flexibility but rather social specialisation. Primates are well known for their flexibility but have little social flexibility. This sounds contradictory, but means that their social behaviour shown towards other group members is flexible, but not the social organisation of their group. In primates, social organisation is most often species specific and barely varies within species, even though many species have large distributions across ecologically variable habitats (44, 45). Baboons (*Papio spec.*), for example, always form multi-male, multi-female groups, even though they occur in a wide range of habitats (45) and gibbons (*Hylobates spec.*) always live in monogamous family groups (45). Primates typically have a clear but flexible dominance hierarchy. Flexible social tactics enable them to respond to arising conflicts, environmental and social changes, stabilizing the social organisation of the group. Thus, flexibility in social behaviour enables primates to maintain their species-specific social organisation.

In contrast, social flexibility (population level) might arise in species that have little flexibility in social behaviour (individual level) towards other group-members. If conflict arises in these species, they might have to change their social organisation to resolve it. This is evident in the socially flexible African striped mouse (*Rhabdomys pumilio*) which can live solitarily, in single family groups or extended family groups (46). In this species, one can distinguish between non-breeding philopatric males and females, one breeding male and up to four breeding females per group (47). Surprisingly, no measurable dominance hierarchy exists and groups are highly egalitarian (48). If conflict occurs, no changes in dominance hierarchy are possible nor do submissive behaviours exist to resolve conflict. Instead, striped mice switch to a solitary lifestyle if conflict increases (46, 49). Social flexibility might thus arise from the absence of flexibility in social behaviour, making solitary living the only adaptive alternative to group living when inter individual conflict increases to a certain threshold level. In contrast,

species with pronounced dominance hierarchies are able to adjust their social interactions with other individuals (social structure), enabling them to maintain their social organisation.

Social flexibility is predicted to be an adaptation to unpredictably changing environments, selecting for high phenotypic flexibility that is based on a broad reaction norm, not on genetic polymorphisms for specific tactics (15). The environment in which it evolved had to be less predictable than in the case of developmental plasticity, such that the environment in which an individual grows up does not provide significant information about the environment in which the same individual will reproduce. Thus, selection favours individuals that can change their social behaviour as adults. If this occurs in both sexes, the entire social organisation of a population can change. For example, both male and female striped mice can be highly sociable and form extended family groups, but the same individuals can switch to solitary living with very few if any social interactions (15, 46, 49). The fact that groups can exist even under conditions of low population density but split at the beginning of the breeding season (individuals leave the group) indicates that the individuals have a choice between group- and solitary living. Thus, the change in social organisation is a consequence of individual decisions (50) and not entirely due to extrinsic environmental factors (see below).

Social flexibility occurs in several species, including insects, birds and mammals (examples in Tab. 1). The most important factor selecting for social flexibility seems to be unpredictable fluctuations in population density influencing the extent of intra-specific competition. For example, in African striped mice, population density declined from one generation to the next from 32.4 to 1.5 mice/ha (factor of 21) due to an unexpected drought, and only a few generations (=years) later due to an unexpected and local increase of predation pressure (presence of a single wild cat) from 30.5 to 6.5 mice /ha (factor of 5) (46). In both cases, striped mice grew up under high population densities, when group-living and communal breeding were the best option, but they reproduced when population density was low and thus solitary breeding was the best option. Under high population density, inter-group competition

in the form of territoriality is very strong (51) leading to small territories (52) and constrained dispersal, such that breeding in groups is favoured (46). However, living in groups can lead to female-female aggression and female infanticide (46) as well as sexual suppression of males by the dominant male (53), resulting in significant fitness costs that can be avoided by becoming solitary (49). Thus, under low population density, individual costs of inter-group competition are lower than costs of intra-group competition, making the switch from group- to solitary-living adaptive.

4. ENTIRELY EXTRINSIC FACTORS

Intra-specific variation in social organisation can also result solely from extrinsic factors (stochastic processes) leading to non-adaptive changes in social organisation. As such there can be no ultimate or proximate explanation for it. In this case the observed intra-specific variation in social organization is a direct consequence of a demographic interruption, and this variation is not due to an adaptive response of individuals to environmental change. Hereby the individuals will be forced to show flexibility in their behaviour to respond adaptively to the change (such as starting mate searching, dispersal), but this flexibility is not the reason for the observed intra-specific variation in social organisation that we want to explain but its consequence (Fig. 1).

In socially monogamous species, the natural death of one of the dominant breeders will change the social organisation not because the remaining family members chose this new social organisation, but simply because the disappearance of one breeder changes group composition. For example, Callitrichids, small New World monkeys, have been reported to show the highest degree of social flexibility in primates (45, 54), but in several species this might be due to social disruption (mortality of breeder) of the default social organisation of pair-living (12). Nevertheless, if such disruption occurs regularly and has significant fitness effects (55-57) it could function as an important selection pressure for social flexibility. On the

other hand, the significant fitness costs associated with a deviation from social monogamy, especially female infanticide (55, 56), might explain the absence of social flexibility and instead a disposition to re-arrange social monogamy after social disruption. Thus, instead of the evolution of social flexibility, evolution of flexibility in social behaviour that promotes a return to a socially monogamous situation might have been favoured.

Individuals may be constrained to live solitarily when population density is very low, and constrained to be group-living when population density is very high, without giving individuals the *choice* to remain in a group or to become solitary. In some species, it is well known that the social organisation is rather inflexible and not influenced by population density. Obligate social species form groups even under very low population densities, for example lions (*Panthera leo*) in the Kalahari (58). On the other hand, some species like whistling rats (*Parotomys brantsii*) live solitarily even under very high population densities (59). Thus, while ecological factors are important in explaining intra-specific variation in social organisation, it is not clear whether these extrinsic factors alone are sufficient to explain differences between species.

5. PHYSIOLOGICAL MECHANISMS UNDERLYING INTRA-SPECIFIC VARIATION IN SOCIAL ORGANISATION

We know that the gene-environment interaction determines phenotypes (including behavioural phenotypes). Thus, changes in both genotypes and in the environment can induce changes in social behaviours. The four different processes that can explain intra-specific variation in social organisation are predicted to have evolved in different environments with different selection pressures (Tab. 2). The underlying physiological factors, themselves being the result of evolution, differ accordingly (Tab. 2). Here the concept of organisational vs. activational effects is important (60). Organisational effects occur early in development and are non-reversible, as in the case of sex determination in mammals (60). Activational effects typically

occur in adulthood and are reversible, for example the activation of sexual behaviour by testosterone (60).

In species where genetic variation explains the variation in social organisation, two or more genotypes occur that cause different developmental pathways characterized by different organisational effects, such as in the case of genetically determined ARTs (36). In the case of developmental plasticity only one genotype occurs, but depending on the environment this can lead to alternative developmental pathways that will rely on different organisational effects, resulting in a determined, non-flexible adult phenotype (35).

Physiological mechanisms of social flexibility are expected to consist of activational, reversible effects such as environmentally induced neural activation or secretion of specific hormones or neuropeptides. The effects of many endocrine parameters on changes in social behaviour have been demonstrated, such as those of prolactin and steroid hormones on parental care (61), and those of testosterone on reproductive behaviour (62). Neuroendocrine changes, e.g. in the production and secretion of neuropeptides (oxytocin, vasopressin), have been demonstrated to be important too, but have been shown only in a few species because of the difficulties of assessing neuropeptides (63). Finally, activation of existing neuronal pathways and neural mechanisms of learning can be expected to be important, but are difficult to study and poorly understood. In sum, social flexibility is regulated by activational physiological mechanisms, enabling an adaptive and reversible response to unpredictable environments.

Genetically or developmentally determined neuronal pathways can be superior to flexible ones, because flexibility requires energy, time, and the opportunity to change (64). In the neurosciences, it is widely held that genetically fixed neuronal patterns are favoured under long-term environmental stability, but that flexibility will be favoured when there is significant environmental instability (65). Developmental plasticity is more advantageous than genetic determinism when change occurs predictably. Flexibility, by learning or by physiological activational effects, is favoured over genetically or developmentally determined pathways

when change happens unpredictably. This is related to the Baldwin effect which states that while acquired traits cannot be inherited, the tendency to acquire traits can be inherited (66). Thus, the Baldwin effect describes the evolution of the ability to respond optimally to a particular environment. This is the result of selection for genes for plasticity and flexibility enabling adaptation to the current environment, rather than genes for a fixed phenotype (67). However, while the idea of the Baldwin effect is more than 100 years old and many studies have demonstrated how organisms rapidly respond both physiologically and behaviourally to changing environments, we still cannot easily link functional importance and inheritance of novel accommodations, i.e. the idea that natural selection sorts among developmental variants for genes enabling plasticity and flexibility (68). In other words: we do not know whether flexible genes *vs.* inflexible genes exist (but see (69)).

In the case of extrinsic factors alone causing intra-specific variation in social organisation, no specific physiological mechanisms exist that lead to the change of social organisation, which entirely depends on the environment. Instead, a stress response will be activated, leading to behavioural changes to return to the default social organisation (70).

6. FUTURE RESEARCH

While the four suggested processes offer an approach to study ultimate and proximate explanations for intra-specific variation in social organisation, many unanswered questions remain:

1. What are the benefits of having a narrow reaction norm (environmental canalization reducing environmental influence on the phenotype) versus having a broad reaction norm? Constraints for the evolution of phenotypic plasticity have received significant theoretical attention, and many studies addressed the costs (same phenotype of plastic genotype has lower fitness than of specialised genotype) and limits (plastic phenotype cannot produce the trait as well as the specialised genotype) of phenotypic plasticity (34,

71, 72). However, decades of research revealed that costs of phenotypic plasticity are surprisingly low (71, 73) and we still do not understand why so many species are specialists instead of being plastic generalists. Few if any empirical studies directly compared specialised *vs.* plastic species, asking under which circumstances can flexible species outcompete specialised ones and vice versa.

2. What environmental selection pressures lead to the evolution of developmental plasticity, which ones to the evolution of social flexibility? Theoretical models exist to explain the evolution of plasticity (developmental plasticity or flexibility) (34) as well as its costs, limits and constraints (71). If the current environment reliably predicts the future environment, developmental plasticity is predicted to evolve, but with increasing unpredictability social flexibility might evolve (Tab. 2), enabling a quick response that can be reversed if the environment changes again. This might be especially important in (1) highly unpredictable environments, and (2) in long-lived species, as predictability tends to decrease with increasing time in the future. In contrast, developmental plasticity resulting in an inflexible phenotype might reduce specific costs of plasticity, especially maintenance and information-acquisition costs: for developmental plasticity, individuals only need to acquire information about the environment once in early development, when deciding which developmental trajectory to follow.

All four processes mentioned here are possible explanations for intra-specific variation in social organisation, but few studies have identified which one explains the pattern found in a specific population or species. In Tab. 3 I provide several predictions for the four different processes that can be used to determine which one explains the pattern of intra-specific variation in social organisation observed in a particular species. While writing this review, it became evident to me how little we know about the processes underlying intra-specific variation in social organisation. I wanted to include many empirical examples for the four possible processes, but apart from my own research topic – social flexibility – not much

literature exists for the other processes. Of course, flexibility might be the most common process for intra-specific variation in social organisation and the other processes might be less important. Genetic variation and developmental plasticity might be important in the expression of individual differences in social behaviour (for example ARTs), but rarely explain intra-specific variation in social organisation. Entirely extrinsic factors might be the most overlooked process, because even in this case ecological factors do exist (stochastic mortality by a predator is ultimately also due to an ecological factor), and researchers in behavioural ecology are prone to see intra-specific variation as an adaptive response to a changing environment, without taking non-adaptive alternatives into account.

Environmental factors are important regulators of intra-specific variation in social organisation, and these factors have been studied extensively (1, 15, 17, 74, 75). For all four processes, the environment is a significant factor, either as a selection pressure (genetic variation), because the current environment determines the best tactic for the next generation but not for more distant generations (developmental plasticity), or because only the current environment determines the best tactic for the currently breeding individuals (social flexibility). Thus, to really understand both the ultimate reasons and the proximate causes of intra-specific variation in social organisation, we have to go further than focussing on the environmental factors. By a combination of only two studies one can differentiate between the four processes (Tab. 4).

The resilience of a species, population or an individual to environmental change depends on its ability to respond adaptively. To understand how species are able to adapt to changing environments we need to understand the different mechanisms of intra-specific variation in social organisation, which is the consequence of individual adaptive responses. If environmental change happens slowly, fluctuates predictably or if environments differ between populations, then genetic variation might be the evolutionary most stable response. If changes happen faster than genetic adaptation, phenotypic plasticity or flexibility can enable an

adaptive response. Developmental plasticity provides animals with opportunity to mount a response during their growth phase that is adaptive when they reproduce later as adults. Social flexibility results from the immediate response of individuals to current changing environmental conditions. In contrast to developmental plasticity, social flexibility is reversible and evolves in highly unpredictable environments. In highly predictable environments, plasticity and flexibility are not favoured by natural selection and instead specialisation evolves. Understanding these different processes of intra-specific variation in social organisation will help us to understand the evolution of sociality and adaptability of species.

Important comments by Loren Hayes, Neville Pillay, Nancy Solomon, Josh van Buskirk and three referees significantly improved this manuscript.

References

1. Krebs JR, Davies NB. 1993 *An introduction to behavioural ecology*. 3rd ed. Oxford, UK: Blackwell Science Ltd.
2. Crook JH. 1964 Evolution of social organisation and visual communication in the weaver birds (Ploceinae). *Behaviour Suppl.* **10**, 1-178
3. Jarman PJ. 1974 The social-organization of antelope in relation to their ecology. *Behaviour.* **48**, 215-267
4. Goldizen W. 1990. A comparative perspective on the evolution of tamarin and marmoset social systems. *Int. J. Primatol.* **11**, 63-83
5. Hayes LD. 2000 To nest communally or not to nest communally: a review of rodent communal nesting and nursing. *Anim. Behav.* **59**, 677-88
6. Kappeler PM, Schaik CPv. 2002 Evolution of primate social systems. *Int. J. Primatol.* **23**, 23:707-40.
7. Lott DF. 1991 *Intraspecific variation in the social systems of wild vertebrates*. New York, USA: Cambridge University Press
8. Lott DF. 1984 Intraspecific variation in the social systems of wild vertebrates. *Behaviour.* **88**, 266-325
9. Kamilar JM. 2013 Phylogenetic approaches to understanding primate behavioural diversity. *Philos. Trans. R. Soc. B.* XXX
10. Clutton-Brock TH, Lukas D. 2013 Evolutionary transitions and constraints on adaptation. *Philos. Trans. R. Soc. B.* XXX
11. Thierry B. 2013 External and internal constraints in socioecology. *Philos. Trans. R. Soc. B.* XXX
12. Anzenberger G, Falk B. 2012 Monogamy and family life in callitrichid monkeys: deviations, social dynamics and captive management. *Int. Zoo Yearbook.* **46**, 109-122

13. Silk JB. 2007. The adaptive value of sociality in mammalian groups. *Philos. Trans. R. Soc. B.* **362**, 539-559
14. Brashares JS, Garland T, Arcese P. 2000 Phylogenetic analysis of coadaptation in behavior, diet, and body size in the African antelope. *Behav. Ecol.* **4**, 452-463
15. Schradin C, Lindholm AK, Johannesen J, Schoepf I, Yuen C-H, König B, Pillay N. 2012 Social flexibility and social evolution in mammals: a case study of the African striped mouse (*Rhabdomys pumilio*). *Molec. Ecol.* **21**, 541-553
16. Maher CR, Burger JR. 2011 Intraspecific variation in space use, group size, and mating systems of caviomorph rodents. *J. Mammal.* **92**, 54-64
17. Koenig WD, Pitelka FA, Carmen WJ, Mumme RL, Stanback MT. 1992 The evolution of delayed dispersal in cooperative breeders. *Quart. Rev. Biol.* **67**, 111-150
18. Stirling DG, Réale D, Roff DA. 2002 Selection, structure and the heritability of behaviour. *J. Evol. Biol.* **15**, 277-289
19. Anholt RRH, Mackay TFC. 2010 *Genetics of social interactions. Principles of Behavioral Genetics*. Amsterdam: Elsevier. p. 185-206
20. Rymer T, Pillay N. 2011 Transmission of parental care behavior in African striped mice, *Rhabdomys pumilio*. *J. Exp. Zool A: Ecol. Gen. Physiol.* **315**, 631-638
21. Hammock EAD, Young LJ. 2005 Microsatellite instability generates diversity in brain and sociobehavioral traits. *Science*. **308**, 1630-1634
22. Maynard Smith J. 1977 Parental investment - prospective analysis. *Anim. Behav.* **25**, 1-9
23. Schradin C, Pillay N. 2005 The influence of the father on offspring development in the striped mouse. *Behav. Ecol.* **16**, 450-455
24. Cushing BS, Razzoli M, Murphy AZ, Epperson PM, Le W, Hoffman GE. 2004 Intraspecific variation in estrogen receptor alpha and the expression of male sociosexual behavior in two populations of prairie voles. *Brain Res.* **1016**, 247- 254
25. Ophir AG, Phelps SM, Sorin AB, Wolff JO. 2007 Morphological, genetic and behavioral comparisons of two prairie vole populations in the field and laboratory. *J. Mammal.* **88**, 989-99
26. Maynard Smith J. 1974 The theory of games and the evolution of animal conflicts. *J. Theor. Biol.* **47**, 209-221
27. Sinervo B, Zamudio KR. 2001 The evolution of alternative reproductive strategies: fitness differential, heritability, and genetic correlation between the sexes. *J. Hered.* **92**, 198-205.
28. Taborsky M. 2008 Alternative reproductive tactics in fish. In: *Alternative reproductive tactics* (eds. RF Oliveira, M Taborsky, HJ Brockmann). pp. 251-299. Cambridge, UK: Cambridge University Press
29. Lank DB, Smith CM, Hanotte O, Burke T, Cooke F. 1995 Genetic polymorphism for alternative mating behaviour in lekking male ruff *Philomachus pugnax*. *Nature*. **378**, 59-62
30. Shuster SM, Sassaman C. 1997 Genetic interaction between male mating strategy and sex ratio in a marine isopod. *Nature*. **388**, 373-376
31. Gotzek D, Ross KG. 2007 Genetic regulation of colony social organization in fire ants: An integrative overview. *Quart. Rev. Biol.* **82**, 201-226
32. Simpson GG. 1953 The Baldwin effect. *Evolution*. **7**, 110-117
33. Bradshaw AD. 1965 Evolutionary significance of phenotypic plasticity in plants. In: *Advances in Genetics* (eds. EW Caspari, JM Thoday) pp. 115-155. Academic Press
34. Chalambor CK, Angeloni LM, Carroll SP. 2010 Behavior as phenotypic plasticity. In: *Evolutionary Behavioral Ecology* (eds. DF Westneat, CW Fox), pp. 90-107. Oxford, UK: Oxford University Press
35. Piersma T, Drent J. 2003 Phenotypic flexibility and the evolution of organismal design. *TREE*. **18**, 228-233

36. Moore MC, Hews DK, Knapp R. 1998 Hormonal control and evolution of alternative male phenotypes: generalizations of models for sexual differentiation. *Amer. Zool.* **38**, 133-151
37. Ellers J, Stuefer J. 2010 Frontiers in phenotypic plasticity research: new questions about mechanisms, induced responses and ecological impacts. *Evol. Ecol.* **24**, 523-526
38. vom Saal F, Bronson F. 1980 Sexual characteristics of adult female mice are correlated with their blood testosterone levels during prenatal development. *Science*. **208**, 597-599
39. Weaver ICG, Cervoni N, Champagne FA, D'Alessio AC, Sharma S, Seckl JR, Dymov S, Szyf M, Meaney MJ. 2004 Epigenetic programming by maternal behavior. *Nature Neurosc.* **7**, 847-854
40. Siegeler K, Sachser N, Kaiser S. 2011 The social environment during pregnancy and lactation shapes the behavioral and hormonal profile of male offspring in wild cavies. *Develop. Psychobiol.* **53**, 575-584
41. Heiming RS, Bodden C, Jansen F, Lewejohann L, Kaiser S, Lesch K-P, Palme R, Sachser N. 2011 Living in a dangerous world decreases maternal care: A study in serotonin transporter knockout mice. *Horm Behav.* **60**, 397-407
42. Champagne FA. 2012 Epigenetics and developmental plasticity across species. *Develop. Psychobiol.* (doi:10.1002/dev.21036)
43. Schradin C, Lindholm AK. 2011 Relative fitness of alternative male reproductive tactics in a mammal varies between years. *J. Anim. Ecol.* **80**, 908-917
44. Thierry B, Iwaniuk AN, Pellis SM. 2000 The influence of phylogeny on the social behaviour of macaques (Primates: Cercopithecidae, genus *Macaca*). *Ethology*. **106**, 713-728
45. Shultz S, Opie C, Atkinson QD. 2011 Stepwise evolution of stable sociality in primates. *Nature*. **479**, 219-222
46. Schradin C, König B, Pillay N. 2010 Reproductive competition favours solitary living while ecological constraints impose group-living in African striped mice. *J. Anim. Ecol.* **79**, 15-21
47. Schradin C, Pillay N. 2004 The striped mouse (*Rhabdomys pumilio*) from the succulent karoo of South Africa: A territorial group living solitary forager with communal breeding and helpers at the nest. *J. Comp. Psychol.* **118**, 37-47
48. Schubert M, Pillay N, Schradin C. 2009 Parental and allo-parental care in a polygynous mammal. *J. Mammal.* **90**, 724-731
49. Schoepf I, Schradin C. 2012 Better off alone! Reproductive competition and ecological constraints determine sociality in the African striped mouse (*Rhabdomys pumilio*). *J. Anim. Ecol.* **81**, 649-656
50. Schradin C, Schubert M, Pillay N. 2006 Winter huddling groups in the striped mouse. *Can. J. Zool.* **117**, 317-324
51. Schradin C. 2004 Territorial defense in a group living solitary forager: who, where, against whom? *Behav. Ecol. Sociobiol.* **55**, 439-446
52. Schradin C, Pillay N. 2006 Female striped mice (*Rhabdomys pumilio*) change their home ranges in response to seasonal variation in food availability. *Behav. Ecol.* **17**, 452-458
53. Schradin C, Schneider C, Yuen CH. 2009 Age at puberty in male African striped mice: the impact of food, population density and the presence of the father. *Funct. Ecol.* **23**, 1004-1013
54. Garber PA. 1997 One for all and breeding for one: cooperation and competition as a tamarin reproductive strategy. *Evol. Anthropol.* **3**, 187-199
55. Bezerra BM, Souto ADS, Schiel N. 2007 Infanticide and cannibalism in a free-ranging plurally breeding group of common marmosets (*Callithrix jacchus*). *Am. J. Primatol.* **69**, 945-952
56. Digby LJ. 1995 Infant care, infanticide, and female reproductive strategies in polygynous groups of common marmosets (*Callithrix jacchus*). *Behav. Ecol. Sociobiol.* **37**, 51-61

57. Huck M, Löttker P, Böhle UR, Heymann EW. 2004 Paternity and kinship patterns in polyandrous moustached tamarins (*Saguinus mystax*). *Am. J. Phys. Anthropol.* **127**, 449-464.
58. Bothma JdP, Walker C. 1999 *Larger carnivores of the African savannas*. Pretoria, South Africa: van Schaik Publishers;
59. Jackson TP. 1999 The social organisation and breeding system of Brants' whistling rat (*Parotomys brantsii*). *J. Zool. Lond.* **247**, 323-331
60. Phoenix CH, Goy RW, Gerall AA, Young WC. 1959 Organizing action of prenatally administered testosterone propionate on the tissue mediating mating behavior in the female guinea pig. *Endocrinology.* **65**, 369-82
61. Schradin C, Anzenberger G. 1999 Prolactin, the hormone of paternity. *News Physiol. Sci.* **14**, 223-231
62. Wingfield JC. 2009 Hormone-behavior interrelationships in a changing environment. In: *Endocrinology of social relationships* (eds. PT Ellison, PB Gray), pp 74-94. Cambridge, USA: Harvard University Press
63. Porges SW, Carter CS. 2011 Mechanisms, mediators, and adaptive consequences of caregiving. In: *Perspectives from evolutionary biology, neuroscience, and the social sciences: moving beyond self-interest* (eds SL Brown, RM Brown, LA Penner), pp 53-74. Oxford, USA: Oxford University Press
64. Gazzinga MS. 2011 Who's in charge? New York, USA: HarperCollins Publisher
65. Papineau D. 2005 Social learning and the Baldwin effect. In: *Evolution, rationality and cognition: A cognitive science for the twenty-first century*. (ed A Zilao), pp. 40-60. New York, USA: Routledge
66. Baldwin JM. 1896 A new factor in evolution. *Amer. Nat.* **30**, 441-451
67. Krubitzer L, Kaas L. 2005 The evolution of the neocortex in mammals: how is phenotypic diversity generated? *Cur. Op. Neurobiol.* **15**, 444-453
68. Badyaev AV. 2009 Evolutionary significance of phenotypic accommodation in novel environments: an empirical test of the Baldwin effect. *Philos. Trans. Roy. Soc. B.* **364**, 1125-1141
69. Silander OK, Nikolic N, Zaslaver A, Bren A, Kikoin I, Alon U, Ackermann M. 2012. A genome-wide analysis of promoter-mediated phenotypic noise in *Escherichia coli*. *PloS Genetics.* **8**, e1002443
70. Mendoza SP. 1991 Behavioural and physiological indices of social relationships: comparative studies of New World monkeys. In: *Primate responses to environmental change* (ed HO Box), pp. 311-335. London, UK: Chapman and Hall
71. Auld JR, Agrawal AA, Relyea RA. 2010 Re-evaluating the costs and limits of adaptive phenotypic plasticity. *Proc R Soc B-Biol Sci.* **277**, 503-511
72. Givnish TJ. 2002 Ecological constraints on the evolution of plasticity in plants. *Evol. Ecol.* **16**, 213-242
73. VanBuskirk J, Steiner UK. 2009 The fitness costs of developmental canalization and plasticity. *J. Evol. Biol.* **22**, 852- 860
74. Lacey EA, Wieczorek JR. 2003 Ecology of sociality in rodents: a Ctenomyid perspective. *J. Mammal.* **84**, 1198-1211
75. Solomon NG, Keane B. 2012 Make space enough between you: Intraspecific variation in animal spacing. In: *Animal behavior 3 vols* (ed K Yasukawa): Praeger, Connecticut, USA
76. Eggert A-K. 1992 Alternative male mate-finding tactics in burying beetles. *Behav. Ecol.* **3**, 243-254
77. Müller JF, Braunisch V, Hwang W, Eggert A-K. 2006 Alternative tactics and individual reproductive success in natural associations of the burying beetle, *Nicrophorus vespilloides*. *Behav. Ecol.* **18**, 196-203
78. Eggert A-K, Müller JK. 2000 Timing of oviposition and reproductive skew in cobreeding female burying beetles (*Nicrophorus vespilloides*). *Behav. Ecol.* **11**, 357-366

79. Reyer HU. 1980 Flexible helper structure as an ecological adaptation in the pied kingfisher (*Ceryle rudis rudis* L.). *Behav. Ecol. Sociobiol.* **6**, 219-227
80. Reyer H-U. 1984 Investment and relatedness: a cost/benefit analysis of breeding and helping in the pied kingfisher (*Ceryle rudis*). *Anim. Behav.* **32**, 1163-1178
81. McGuire B, Getz LL. 1998 The nature and frequency of social interactions among free-living prairie voles (*Microtus ochrogaster*). *Behav. Ecol. Sociobiol.* **43**, 271-279
82. Lucia KE, Keane B, Hayes LD, Lin YK, Schaefer RL, Solomon NG. 2008 Philopatry in prairie voles: an evaluation of the habitat saturation hypothesis. *Behav. Ecol.* **19**, 774- 783
83. Ophir AG, Phelps SM, Sorin AB, Wolff JO. 2008 Social but not genetic monogamy is associated with greater breeding success in prairie voles. *Anim. Behav.* **75**, 1143-1154
84. Solomon NG, Crist TO. 2008 Estimates of reproductive success for group-living prairie voles, *Microtus ochrogaster*, in high-density populations. *Anim. Behav.* **76**, 881-892
85. Berry RJ, Tattersall FH, Hurst J. 2008 Genus *Mus*. In: *Mammals of the British Isles Handbook 4th ed* (eds S Harris, DW Yalden), Southampton, U.K.: The Mammal Society
86. Latham N, Mason G. 2004 From house mouse to mouse house: the behavioural biology of free-living *Mus musculus* and its implications in the laboratory. *Appl. Anim. Behav. Sci.* **86**, 261-289




Figure 1. The four processes that can lead to intra-specific variation in social organisation. To explain the variation observed at the population level, we need to consider variation at the individual level. Only the two processes on the right represent phenotypic plasticity.

Top row: different individuals differ in social behaviour. Left, genetic variation: individuals of the same species but two different populations differ genetically, leading to differences in social organisation. Right, developmental plasticity: depending on the environment in which individuals grow up, environmental cues will activate developmental pathways for social behaviour that either leads to the same social organisation as observed in their parent generation (far right bottom) or to a different kind of social organisation (far right top).

Bottom row: the same individuals occur in different forms of social organisation. Left, entirely extrinsic factors: the unpredictable death / disappearance of some breeding individuals is the only cause of changes in social organisation. This itself will cause behavioural flexibility in the survivors, such as mate searching, to return to the original state, but flexibility is the result, not the cause of the observed intra-specific variation in social organisation. Right, social flexibility: if the environment changes, social tactics of individuals change, which as a consequence will change the social organisation of the population. This is reversible, as the same individuals can switch their tactic again, if the environment changes again.

Table 1. Species showing social flexibility.

Species	Male tactics	Female tactics	Social organisations	Fluctuation in population density?	Importance of intra-specific competition	References
Burying beetle <i>Nicrophorus vespilloides</i>	<ul style="list-style-type: none"> Attract females via pheromones Attract females to carcass Satellite 	<ul style="list-style-type: none"> Single breeding Communal breeding Parasite 	<ul style="list-style-type: none"> Solitary Pair-living One male, multi female 	Yes	Carion size determines number of females that can breed together.	(76-78)
Pied kingfisher <i>Ceryle rudis</i>	<ul style="list-style-type: none"> Breeder Non-reproducing helper Reproducing helper 	<ul style="list-style-type: none"> Breeder 	<ul style="list-style-type: none"> Pair-living Family group Two males, one female 	Yes	Availability of good nesting sites influences whether un-related helpers are accepted.	(79, 80)
Prairie vole <i>Microtus ochrogaster</i>	<ul style="list-style-type: none"> Philopatric helper Solitary wanderer Breeder 	<ul style="list-style-type: none"> Philopatric helper Single breeder Communal breeder 	<ul style="list-style-type: none"> Solitary Pair-living Multi male, multi female 	Yes (breeding seasons vs. winter)	Female-female competition in communally breeding groups: higher reproductive success if only one female breeds per group.	(81-84)

House mouse <i>Mus musculus</i>	<ul style="list-style-type: none"> • Philopatric helper • Solitary roamer • Breeder 	<ul style="list-style-type: none"> • Philopatric helper • Single breeder • Communal breeder 	<ul style="list-style-type: none"> • Solitary • Pair-living • Multi male, multi female 	Yes	Intra-sexual aggression. Female infanticide.	(85, 86)
Striped mouse <i>Rhabdomys pumilio</i>	<ul style="list-style-type: none"> • Philopatric helper • Solitary roamer • Breeder 	<ul style="list-style-type: none"> • Philopatric helper • Single breeder • Communal breeder 	<ul style="list-style-type: none"> • Solitary • Pair-living • Multi male, multi female 	Yes	Intra-sexual aggression. Female infanticide Male reproductive suppression.	(46, 47, 49, 53)

Table 2. Comparison of the four processes that can cause intra-specific variation in social organisation.

	Genetics	Influence of early environment on behaviour²	Variability within individuals?	Environment in which it evolved	Physiological mechanisms
1. Genetic variation	Polymorphism Narrow reaction norm	No	No	Predictable	Organisational
2. Developmental plasticity	Monomorphism ¹ Broad reaction norm	Non-reversible	No	Short term: predictable Long term: unpredictable	Organisational
3. Social flexibility	Monomorphism ¹ Broad reaction norm	Reversible	Yes	Unpredictable	Activational
4. Entirely extrinsic factors	Monomorphism ¹ Narrow reaction norm	No	Yes	Predictable	None that leads to variation in social organisation

¹The term genetic monomorphism does not imply that genetic variation is absent, only that the major part of variation observed in social behaviour is not due to genetic but to environmental factors.

²Other behaviours can be influenced during early environment, but this cannot explain the variation in social organisation.

Table 3. The predictions to differentiate between the four processes that bring about intra-specific variation in social organisation.

Study	Trait investigated	Yes (trait exists)	No (trait does not exist)
1	Variability within individuals?	Social flexibility or extrinsic factors	Genetic variation or developmental plasticity
2	Genetic polymorphism?	Genetic variation	Developmental plasticity, social flexibility or extrinsic factors
3	Early environment can induce changes in individual behaviour	Developmental plasticity or social flexibility	Genetic variation, extrinsic factors
4	Alternative forms of social organisation are stable	Genetic variation, developmental plasticity, or social flexibility	Extrinsic factors
5	Physiological mechanisms organisational?	Genetic variation or developmental plasticity	Social flexibility
6	Predictability of environment in which it evolved	This factor can change significantly between generations (thus between years, decades or centuries) and is difficult to use for categorisation. Its best used for comparative studies.	

Table 4. Key to determine the process leading to intra-specific variation in social organisation, based on the predictions from Tab. 3. Predictions 5 from Tab. 3 should be used to confirm the results obtained from this key.

No	Question	Result	Go to
1	a Variability occurs within individuals b Variability does not occur within individuals		3 2
2	a Genetic polymorphism b Genetic monomorphism	Genetic variation Developmental plasticity	
3	a Early environment induces changes in individual behaviour and alternative forms of social organisation are stable b Early environment induces no changes in individual behaviour and alternative forms of social organisation are not stable	Social flexibility Entirely extrinsic factors	